

Animating Rotation with Quaternion Curves

Ken Shoemake†

The Singer Company
Link Flight Simulation Division

ABSTRACT

Solid bodies roll and tumble through space. In computer animation, so do cameras. The rotations of these objects are best described using a four coordinate system, quaternions, as is shown in this paper. Of all quaternions, those on the unit sphere are most suitable for animation, but the question of how to construct curves on spheres has not been much explored. This paper gives one answer by presenting a new kind of spline curve, created on a sphere, suitable for smoothly in-betweening (i.e. interpolating) sequences of arbitrary rotations. Both theory and experiment show that the motion generated is smooth and natural, without quirks found in earlier methods.

C.R. Classification: G.1.1 [Numerical Analysis] Interpolation—Spline and piecewise polynomial interpolation; G.1.2 [Numerical Analysis] Approximation—Spline and piecewise polynomial approximation; I.2.9 [Artificial Intelligence] Robotics—Manipulators; I.3.5 [Computer Graphics] Computational Geometry and Object Modelling—Curve, surface, solid, and object representation, —Geometric algorithms, languages, and systems, —Hierarchy and geometric transformations

General Terms: Algorithms, Theory

Keywords and phrases: quaternion, rotation, spherical geometry, spline, Bézier curve, B-spline, animation, interpolation, approximation, in-betweening

1. Introduction

Computer animation of three dimensional objects imitates the *key frame* techniques of traditional animation, using key positions in space instead of key

Permission to copy without fee all or part of this material is granted provided that the copies are not made or distributed for direct commercial advantage, the ACM copyright notice and the title of the publication and its date appear, and notice is given that copying is by permission of the Association for Computing Machinery. To copy otherwise, or to republish, requires a fee and/or specific permission.

drawings. Physics says that the general position of a rigid body can be given by combining a translation with a rotation. Computer animators key such transformations to control both simulated cameras and objects to be rendered. In following such an approach, one is naturally led to ask: What is the best representation for general rotations, and how does one in-between them? Surprisingly little has been published on these topics, and the answers are not trivial.

This paper suggests that the common solution, using three Euler's angles interpolated independently, is not ideal. The more recent (1843) notation of quaternions is proposed instead, along with interpolation on the quaternion unit sphere. Although quaternions are less familiar, conversion to quaternions and generation of in-between frames can be completely automatic, no matter how key frames were originally specified, so users don't need to know—or care—about inner details. The same cannot be said for Euler's angles, which are more difficult to use.

Spherical interpolation itself can be used for purposes besides animating rotations. For example, the set of all possible directions in space forms a sphere, the so-called Gaussian sphere, on which one might want to control the positions of infinitely distant light sources. Modelling features on a globe is another possible application.

It is simple to use and to program the method proposed here. It is more difficult to follow its development. This stems from two causes: 1) rotations in space are more confusing than one might think, and 2) interpolating on a sphere is trickier than interpolating in, say, a plane. Readers well acquainted with splines and their use in computer animation should have little difficulty, although even they may stumble a bit over quaternions.

2. Describing rotations

2.1 Rigid motion

Imagine hurling a brick towards a plate glass window. As the brick flies closer and closer, a nearby physicist



might observe that, while it does not change shape or size, it can tumble freely. Leonhard Euler proved two centuries ago that, however the brick tumbles, each position can be achieved by a single rotation from a reference position. [Euler,1752] [Goldstein] The same is true for any rigid body. (Shattering glass is obviously not a single rigid body.)

While translations are well animated by using vectors, rotation animation can be improved by using the progenitor of vectors, quaternions. Quaternions were discovered by Sir William Rowan Hamilton in October of 1843. The moment is well recorded, for he considered them his most important contribution, the inspired answer to a fifteen-year search for a successor to complex numbers. [Hamilton] By an odd quirk of mathematics, only systems of two, four, or eight components will multiply as Hamilton desired; triples had been his stumbling block.

Soon after quaternions were introduced, Arthur Cayley published a way to describe rotations using the new multiplication. [Cayley] The notation in his paper so closely anticipates matrix notation, which he devised several years later, that it may be taken as a formula for converting a quaternion to a rotation matrix. It turns out that the four values making up a quaternion describe rotation in a natural way: three of them give the coordinates for the axis of rotation, while the fourth is determined by the angle rotated through. [Courant & Hilbert]

Since computer graphics leans heavily on vector operations, it is perhaps easiest to explain quaternions and rotation matrices in terms of these, reversing history. However quaternions can stand on their own as an elegant algebra of space. [Herstein] [Pickert] [MacLane]

2.2 Rotation matrices

That a tumbling brick does not change size, shape, nor "handedness" is mathematically expressed as the preservation of dot products and cross products, since these measure lengths, angles, and handedness. And since the determinant of a 3×3 matrix can be computed as the dot product of one column with the cross product of the other two, determinants are also preserved. Symbolically:

$$\text{Rot}(u_1) \cdot \text{Rot}(u_2) = u_1 \cdot u_2$$

$$\text{Rot}(u_1) \times \text{Rot}(u_2) = \text{Rot}(u_1 \times u_2)$$

$$\det(\text{Rot}(u_1), \text{Rot}(u_2), \text{Rot}(u_3)) = \det(u_1, u_2, u_3)$$

An immediate consequence is that orientation changes must be linear operations, since the preserved operations are; hence they have a matrix representation, M . Using the matrix form of a dot product, $u_1^t u_2$, we can say more precisely that $(M u_1)^t (M u_2) = u_1^t u_2$, from which it follows that

$$M^t M = I.$$

That is, the change matrix M is *orthogonal*; its columns (and rows) are mutually perpendicular unit magnitude vectors. Because M must also preserve determinants, it is a *special orthogonal* matrix, satisfying

$$\det(M) = +1.$$

It is well known, and anyhow easy to show, that the special orthogonal matrices form a group, $SO(3)$, under multiplication. [MacLane][Goldstein][Misner] In this rotation group, the inverse of M is just M^t , the opposite rotation.

To illustrate, the matrix

$$M = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta \\ 0 & \sin \theta & \cos \theta \end{bmatrix}$$

effects a rotation through an angle of θ around the x axis. After verifying the properties discussed so far, note that the diagonal entries sum to $1 + 2\cos \theta$. While it is too lengthy to show here, the diagonal sum measures the same quantity for matrices generating rotation around any axis. [MacLane]

2.3 Quaternions

Quaternions, like rotations, also form a non-commutative group under their multiplication, and these two groups are closely related. [Goldstein] [Pickert][Misner] In fact, we can substitute quaternion multiplication for rotation matrix multiplication, and do less computing as a result. [Taylor]

To perform quaternion arithmetic, group the four components into a real part—a scalar, and an imaginary part—a vector. Addition is easy: add scalar to scalar and vector to vector. But our major interest is in multiplication. Start with a simple case: multiply two quaternions without real parts, or more precisely, with zero real parts. The result quaternion has a vector that is the cross product of the two vector parts, and a scalar that is their dot product, negated:

$$u_1 u_2 = [(-u_1 \cdot u_2), (u_1 \times u_2)].$$

It is certainly convenient to encompass both vector products with a single quaternion product. (One early lover of quaternion algebra called vector algebra a "hermaphrodite monster", since it required two kinds of product, each yielding a different type of result.) If one quaternion has only a scalar part, with its vector components all zero, multiplication is just real multiplication and vector scaling. Combining the two effects gives the general rule [Brady]:

$$[s_1, u_1] [s_2, u_2] = [(s_1 s_2 - u_1 \cdot u_2), (s_1 u_2 + s_2 u_1 + u_1 \times u_2)].$$

Except for the cross product this looks like complex multiplication, $(a_1 + ib_1)(a_2 + ib_2) = (a_1a_2 - b_1b_2) + i(a_1b_2 + a_2b_1)$, as Hamilton intended.[†]

Quaternions multiply with a cross product because rotations confound axes. To illustrate, place a book in front of you, face up, with the top farthest away. Use this orientation as a reference. Now hold the sides and flip it toward you onto its face, rotating 180 degrees around a left-to-right axis, y . Then, keeping it face down, spin it clockwise 180 degrees around an up-down z axis. Two rotations around two perpendicular axes; yet the total change in orientation must be, according to Euler, a single rotation. Indeed, if you hold the ends of the spine and flip the book 180 degrees around this third, outward-pointing, x axis, you should restore the original orientation. As quaternions, this is—anticipating developments ahead— $[0, (0,1,0)]$ times $[0, (0,0,1)]$ equals $[0, (1,0,0)]$; the cross product is essential.

Notice how quaternion operations give a new orientation, in "quaternion coordinates", much as translations give a position, relative to some starting reference. A central message of this paper is that quaternion coordinates are best for interpolating orientations. For comparison, imagine using spherical coordinates for translations! Quaternions represent orientation as a single rotation, just as rectangular coordinates represent position as a single vector. Translations combine by adding vectors; rotations, by multiplying quaternions. The separate axes of translations don't interact; the axes of rotations must. Quaternions preserve this interdependence naturally; Euler's angle coordinates ignore it.

2.4 Euler's angles

Why, then, do so many animators use Euler's angles? Mostly, I suspect, because quaternions are unfamiliar. Unlike Euler's angles, quaternions are not taught early in standard math and physics curricula. Certainly there is a plethora of arguments against angle coordinates. Euler's angle coordinates specify orientation as a series of three independent rotations about pre-chosen axes. For example, the orientation of an airplane is sometimes given as "yaw" (or "heading") around a vertical axis, followed by "pitch" around a horizontal axis through the wings, followed by "roll" around the nose-to-tail line. These three angles must be used in exactly the order given because rotations do not commute. The ordering of rotation axes used is a matter of convention, as is the particular set of axes, no matter what the order. For instance some physicists use the body centered axes z - x - z , in contrast to the aeronautics z - y - x . At least a dozen different conventions are possible for which series of axes to use. [Kane][Goldstein] The geometry of orientations in Euler's angle coordinates is contorted, and varies with choice of initial coordinate axes. There is no

reasonable way to "multiply" or otherwise combine two rotations. Even converting between rotation matrices and angle coordinates is difficult and expensive, involving arbitrary assumptions and trigonometric functions. In their defense, it must be said that they are handy for solving differential equations—which is how Euler used them. [Euler,1758]

3. In-betweening alternatives

3.1 Straight line in-betweening

It is not immediately obvious how to in-between even two rotation keys. What orientations should an object assume on its journey between them? A natural answer is: take the first key as a reference, and represent the second by describing the single rotation that takes you to it, according to Euler's theorem. The in-between orientations should be positioned along that rotation.

If we plot quaternions as points in four-dimensional space, the straight lines between them give orientations interpolating the end points in exactly the above sense. If we plot Euler's angle coordinates instead, the in-between orientations will try to twist around three different axes simultaneously. This angle interpolation treats the three angles of rotation at each key orientation as a three-dimensional vector whose components are interpolated independently from key to key. Paradoxically, we can not rotate simply except around the special axes chosen for composition. We may even encounter so-called "gimbal lock", the loss of one degree of rotational freedom. Gimbal lock results from trying to ignore the cross product interaction of rotations, which can align two of the three axes. Quaternions are safe from gimbal lock, and so have been used for years to handle spacecraft, where it is unacceptable. [Kane][Mitchell]

3.2 How quaternions rotate

Straight lines between quaternions, however, ignore some of the natural geometry of rotation space. If our interpolated points were evenly spaced along a line, the animated rotation would speed up in the middle. To see why, we must look at how a quaternion converts to a rotation matrix. We rotate a vector by a quaternion so: multiply it on the right by the quaternion and on the left by the inverse of the quaternion, treating the vector as $[0, \mathbf{x}]$.

$$\mathbf{x}' = \text{Rot}(\mathbf{x}) = q^{-1} \mathbf{x} q$$

Though it is not obvious, the result will always be a vector, with a zero scalar component. Notice how this guarantees

$$\text{Rot}(\mathbf{x}_1) \text{Rot}(\mathbf{x}_2) = \text{Rot}(\mathbf{x}_1 \mathbf{x}_2)$$

which implies that dot and cross products are preserved, embedded in the quaternion product.

The inverse of a quaternion is obtained by negating its

[†] Hamilton wrote a quaternion as $s + iv^x + jv^y + kv^z$, with $i^2 = j^2 = k^2 = ijk = -1$. The multiplication rules given before are consequences of this elegant formulation.



vector part and dividing both parts by the magnitude squared. For $q = [s, \mathbf{u}]$,

$$q^{-1} = \frac{1}{||q||^2} [s, -\mathbf{u}]; \quad ||q||^2 = s^2 + \mathbf{u} \cdot \mathbf{u}.$$

Because all effects of magnitude are divided out, any scalar multiple of a quaternion gives the same rotation. (This kind of behavior is not unknown in computer graphics; any scalar multiple of a point in homogeneous coordinates gives the same non-homogeneous point.)

If the scalar part has value w , and the vector part values x , y , and z , the corresponding matrix can be worked out to be

$$M = \begin{bmatrix} 1-2y^2-2z^2 & 2xy+2wz & 2xz-2wy \\ 2xy-2wz & 1-2x^2-2z^2 & 2yz+2wx \\ 2xz+2wy & 2yz-2wx & 1-2x^2-2y^2 \end{bmatrix}$$

when the magnitude $w^2+x^2+y^2+z^2$ equals 1. The magnitude restriction implies that, plotted in four-dimensional space, these quaternions lie on a sphere of radius one. Deeper investigation shows that such unit quaternions carry the amount of rotation in w , as $\cos \theta/2$, while the vector part points along the rotation axis with magnitude $\sin \theta/2$. The axis of a rotation is that line in space which remains unmoved; but notice that's exactly what happens when scalar multiples of \mathbf{u} are rotated by $[s, \mathbf{u}]$. Because the cross product drops out, multiplication commutes, q^{-1} meets q , mutual annihilation occurs, and the vector emerges unscathed. Summing the matrix diagonal leads to the formula stated for w . The sum equals $4w^2-1$, but must also be $1+2\cos \theta$. A trig identity, $\cos 2\theta = 2\cos^2 \theta - 1$, finishes the demonstration.

3.3 Great arc in-betweening

This sphere of unit quaternions forms a sub-group, S^3 , of the quaternion group. Furthermore, the spherical metric of S^3 is the same as the angular metric of $SO(3)$. [Misner] From this it follows that we can rotate without speeding up by interpolating on the sphere. Simply plot the two given orientations on the sphere and draw the great circle arc between them. That arc is the curve where the sphere intersects a plane through the two points and the origin. We sped up before because we were cutting across instead of following the arc; otherwise the paths of rotation are the same.

A formula for spherical linear interpolation from q_1 to q_2 , with parameter u moving from 0 to 1, can be obtained two different ways. From the group structure we find

$$\text{Slerp}(q_1, q_2; u) = q_1(q_1^{-1}q_2)^u;$$

while from the 4-D geometry comes†

$$\text{Slerp}(q_1, q_2; u) = \frac{\sin(1-u)\theta}{\sin \theta} q_1 + \frac{\sin u\theta}{\sin \theta} q_2,$$

where $q_1 \cdot q_2 = \cos \theta$. The first is simpler for analysis, while the second is more practical for applications.

But animations typically have more than two key poses to connect, and here even our spherical elaboration of simple linear interpolation shows flaws. While orientation changes seamlessly, the direction of rotation changes abruptly. In mathematical terms, we want higher order continuity. There are lots of ways to achieve it—off the sphere; unfortunately we've learned too much.

3.4 Rotation geometry and topology


No matter what we do in general quaternion space, the ultimate effect must be interpreted via the sphere; so we had best work there in spite of the difficulty. It is important to grasp this point. The metric structure, hence the intrinsic geometry, of the rotation group $SO(3)$ is that of a sphere. Over small regions, meaning in this case small rotation angles, a sphere looks as if it is flat. But if we go far enough along a "straight line", we end up back where we started. What could be more evident about rotations? Their very essence is moving in circles. Looking back to the book-turning experiment, the confounding of axes is like traveling on a sphere: if we go in some direction to a quarter of the way around the sphere, turn 90 degrees, travel the same distance, then turn and travel again, we will arrive back home, coming in at right angles to the direction we headed out. Even more revealing, we can leave the north pole in any direction and end up at the south pole, just as we can rotate 360 degrees around any axis and end up oriented the same way.

Local geometry does not, however, determine global topology. Contradictory though it may seem, the geometry curves like a sphere, but the topology says north and south poles are the same! In fact, each pair of opposite points represents the same rotation. The reader may preserve sanity through two expedients. One is to see that this, like homogeneous coordinates, is geometry under perspective projection. The second is to restore spherical topology by including "entanglements". Physically, taking an object with strings attached and rotating it 360 degrees leaves the strings tangled; yet—most odd—rotating 720 degrees does not. [Misner][Gardner]

Accepting the topological oddity is more useful here, but it leaves a minor inconvenience. Namely, when converting an orientation in some foreign form, such as a matrix, to a quaternion form, which quaternion should we choose? Which side of the sphere? An answer that works well is this. Construct a string of quaternions through which to interpolate by choosing

† Glenn Davis suggested this formula.

each added quaternion on the side closest to the one before. Then small changes in orientation will yield small displacements on the sphere.



Representing a projective plane

3.5 Splines

We are left with the problem of constructing smooth curves on spheres. About a hundred years after quaternions appeared, Isaac Schoenberg published a two part attack on ballistics and actuarial problems, using what he called splines. [Schoenberg] Named by analogy to a draftsman's tool, these are interpolating curves constructed from cubic polynomial pieces, with second order continuity between pieces. Cubic splines solve an integral equation which says to minimize the total "wobble" of the curve, as measured by the second derivative. These interpolants are very popular, and the equation can be augmented with Lagrange multipliers to constrain the solution curves to lie on a sphere [Courant & Hilbert]; yet there are problems. First, the augmented equation is much more difficult and expensive to solve. Second, the curve must adjust everywhere if one of the points changes; that is, we have no local control.

3.6 Bézier curves

While Schoenberg invented splines based on numerical analysis, Pierre Bézier invented a class of curves, now called by his name, based on geometrical ideas. In fact, he showed how to find points on such a curve by drawing lines and splitting them in regular proportions. [Bézier] This is exactly what is needed. We already know how to do the equivalent—draw great arcs and proportions of arcs—on a sphere. A complete solution needs only a little more.

4. Spherical Bézier curves

4.1 Joining curves


Bézier curves go through only their first and last defining points, but we want to interpolate all our orientations. The trick is to splice together short Bézier curves in the manner of splines. Their creator showed an easy way to do this which guarantees first

order continuity, probably enough for us. As the curve goes through its end points it is tangent to its end segments. Line up the segments across a join, match their lengths, and the curves will piece together smoothly. If the key orientations are placed at joints, then each short curve moves us from one key to the next, because each piece passes through its ends.

Now, although the two segments abutting a curve junction should match each other, one of the segments can be chosen freely. These choices determine the axis and speed of rotation as we pass through the keys. The burden of choice can be passed to the animator of course, but automation is feasible, and generally preferable.

4.2 Choosing joint segments

Spherical linear interpolation gives two conflicting arc segments at a joint, one on each side. Smooth the difference with an even compromise, aiming for a point halfway between where the incoming segment would proceed, and where the outgoing segment must arrive.†



Constructing a point for tangent

Given successive key quaternions q_{n-1} , q_n , q_{n+1} interpreted as 4-D unit vectors, the computation for a segment point a_n after q_n is

$$a_n = \text{Bisect}(\text{Double}(q_{n-1}, q_n), q_{n+1}),$$

where

$$\text{Double}(p, q) = 2(p \cdot q)q - p;$$

$$\text{Bisect}(p, q) = \frac{p + q}{\|p + q\|}.$$


The matching point for the segment before q_n should be

$$b_n = \text{Double}(a_n, q_n)$$

† For the numerically knowledgeable, this construction approximates the derivative at points of a sampled function by averaging the central differences of the sample sequence. [Dahlquist & Björk]



to ensure a smooth join, regardless of how a_n is chosen.



Splicing Bézier segments together


4.3 Evaluating on the sphere

Everything is now in hand to imitate Bézier's curve technique. Each short curve is defined by four quaternions, $q_n, a_n, b_{n+1}, q_{n+1}$. Let the parameter u vary from 0 to 1 as the curve departs q_n towards a_n and arrives at q_{n+1} tangent to the arc from b_{n+1} . Spherically interpolate by proportion u between q_n and a_n , a_n and b_{n+1} , b_{n+1} and q_{n+1} , to obtain three new quaternions. Then interpolate between those to get two more; and finally interpolate again, reducing to a single point. Abbreviating $\text{Slerp}(p, q; u)$ as $(p:q)_u$, the computation looks like this:

$$\begin{aligned} q_n &= p_0^{(0)} \\ (p_0^{(0)}:p_1^{(0)})_u &= p_0^{(1)} \\ a_n &= p_1^{(0)} \quad (p_0^{(1)}:p_1^{(1)})_u = p_0^{(2)} \\ (p_1^{(0)}:p_2^{(0)})_u &= p_1^{(1)} \quad (p_0^{(2)}:p_1^{(2)})_u = p_0^{(3)} = q_{n+u} \\ b_{n+1} &= p_2^{(0)} \quad (p_1^{(1)}:p_2^{(1)})_u = p_1^{(2)} \\ (p_2^{(0)}:p_3^{(0)})_u &= p_2^{(1)} \\ q_{n+1} &= p_3^{(0)} \end{aligned}$$

4.4 Tangents revisited

A simple check proves the curve touches q_n and q_{n+1} at its ends. A rather challenging differentiation shows it is tangent there to the segments determined by a_n and b_{n+1} . However, as with Bézier's original curve, the magnitude of the tangent is three times that of the segment itself. That is, we are spinning three times faster than spherical interpolation along the arc. Fortunately we can correct the speed by merely truncating the end segments to one third their original length, so that a_n is closer to q_n and b_{n+1} closer to q_{n+1} .



Calculating a Bézier curve point recursively

5. Results

5.1 The grand scheme

What have we ended up with? An animator sits at a workstation and interactively establishes a sequence of keys for, say, camera orientation. The interpolating algorithm does not depend on the nature of the interface the animator sees; all needed information is contained in the sequence of keys. Probably the orientations will be represented internally as matrices, so a conversion step follows. The matrices are "lifted" to a sequence of neighboring quaternions, q_n , on the unit sphere. Each quaternion within the sequence will become the endpoint of two spherical Bézier curves. Between each quaternion pair, q_n and q_{n+1} , two additional points, a_n and b_{n+1} , are added to control motion through the joints. At this point, time becomes a parameter along the composite curve. As the frame number increments, the parameter enters and leaves successive curve pieces. Within each piece a local version of the parameter is adjusted to run from 0 to 1. Now the Bézier geometric construction comes into play, producing an interpolated quaternion, q_{n+u} , from q_n , a_n , b_{n+1} , q_{n+1} , and the local parameter, u . Finally the mint-fresh interpolated quaternion is transmuted into a matrix, to be used in rotating a list of object vectors for rendering.

5.2 Properties

A look at one special case is revealing. Suppose all the points to interpolate are spread along a single arc. This means they represent different amounts of rotation around a single axis, in which case quaternion multiplication commutes. Under these special conditions, the formula for the curve sections reduces to

$$q_{n+u} = q_n^{(1-u)^3} a_n^{3(1-u)^2u} b_{n+1}^{3(1-u)u^2} q_{n+1}^{u^3}$$

When this is compared to the standard Bézier polynomial, $p_n(1-u)^3 + a_n 3(1-u)^2u + b_{n+1} 3(1-u)u^2 + q_{n+1}u^3$, it is apparent that addition and multiplication

have become multiplication and exponentiation. Of course, when the points are not on one arc, commutativity fails, so the formula looks much messier.

In the interesting restricted case when the points are spaced evenly and consecutively around an arc, the resulting animation behaves exactly as we would hope: we get smooth, constant speed rotation around the appropriate axis. Notice that we can choose *any* axis for this rotation. This is clearly preferable to interpolation with Euler's angles, where the coordinate axes are special. A more subtle property of *all* quaternion interpolation is that the motion is independent of coordinate axes. So, for example, if we design a move, then rotate the coordinate system arbitrarily, the geometry of the motion will not change. Euler interpolants, unfortunately, will do wildly different things.

5.3 Applicability

Rotations in space are significantly more complicated than rotations in a plane. It is easy to deal with the latter, since only one parameter is involved. Quaternions are out of place in a plane. Joint control in robotics simulations has its own highly specialized body of techniques; and though quaternions have shown up in the literature, they seem less useful in that context. [Brady][Taylor] However, B.K.P. Horn has used a tessellation of the quaternion unit sphere to identify the orientation of an object from its extended Gaussian image; a good reference is [Brou]. Non-rigid motion obviously needs to be handled specially. But for moving a camera eye-point, and for many kinds of object motion, quaternion interpolation has strong advantages.

5.4 Comparisons and complaints

Cost advantages are difficult to estimate. Converting a matrix to a quaternion requires only one square root and three divides plus some adds, at worst. Converting back requires 9 multiplies and 15 adds. While the conversions don't use trigonometric functions, the arc proportioning does. For comparison, angle interpolation requires several trigonometric functions as well as quite a few multiplies and adds to create each interpolated matrix. My experience is that the Bézier scheme is comfortably fast enough for design work, which is the only time speed has mattered. (If, for some application, more speed is essential, non-spherical quaternion splines will undoubtedly be faster than angle interpolation, while still free of axis bias and gimbal lock.)

These interpolants are not perfect, of course. Like all interpolants, they can develop kinks between the interpolated points. There are simple algorithms for adding new sequence points to ordinary splines without altering the original curve [Boehm]; they do not work for this interpolant. And if these curves can be shown to satisfy some variational principal, it will be by chance. It is useful to do this, because any solution to an integral equation like that for splines admits subdivision [Lane et al]; minimum curvature between

end points implies minimum curvature between intermediate points as well. Along these lines, Gabriel and Kajiya, motivated by quaternions, have been developing a technique to find splines on arbitrary Riemannian manifolds by solving differential equations. [Gabriel & Kajiya]

6. Questions

Future research could answer some interesting practical questions. What are these spherical Bézier curves? Is there some abstract characterization of them? Or is there some related interpolant that is well-characterized? In light of the success of the geometric adaptation approach, it appears reasonable to apply the idea to B-splines, which also have a known geometric evaluation technique. [Gordon & Riesenfeld] How do spherical B-splines behave? Is it possible to add new points to a sequence for either kind of curve without disturbing it? How? Can B-splines be made to interpolate, not just approximate, with a simple adjustment of control points? Is there a way to construct a curve parameterized by arc length? This would be very useful. What is the best way to allow varying intervals between sequence points in parameter space? Abandoning the unit sphere, one could work with the four-dimensional Euclidean space of arbitrary quaternions. How do standard interpolation methods applied there behave when mapped back to matrices? Note that we now have little guidance in picking the inverse image for a matrix, and that cusp-free R^4 paths do not always project to cusp-free S^3 paths.

However these questions are answered, quaternion spline interpolants already offer a well-behaved improvement over traditional techniques. They are simple to use, simple to implement, robust, efficient, consistent, and flexible. More research would make them even more so.

7. Acknowledgments

This work was begun for an animation system I designed and implemented at Singer-Link. Several people there deserve thanks, but I especially thank Glenn Davis, who befriended me with his good humor and mathematical efforts as I struggled through trying times.

I prefer not to invent the wheel if I can find the plans; so I pestered Don Venhaus, Brian Barsky, Tom Duff, Lance Williams, and Jim Blinn, whom I thank for their time, their comments, and their assurances that they had not seen this particular wheel roll past PDI, Berkeley, Lucasfilm, NYIT, or JPL.

Thanks to everyone at Pacific Data Images for the interest and encouragement that got me started.

The folks at Ridge Computer were generous above and beyond the call of customer support in letting me use their Imagen typesetting system to produce this paper.

Lastly, I thank Nori Hall for commenting on numerous drafts, and more.

References

1. BÉZIER, P.E., *Numerical Control — Mathematics and Applications*, John Wiley and Sons, London (1972).
2. BOEHM, WOLFGANG, "Inserting new knots into B-spline curves," *Computer-Aided Design* 12(4) pp. 199-201 (July 1980).
3. BRADY, MICHAEL, "Trajectory Planning," in *Robot Motion: Planning and Control*, ed. Michael Brady, John M. Hollerbach, Timothy L. Hohnson, Tomas Lozano-Perez and Matthew T. Mason, The MIT Press (1982).
4. BROU, PHILIPPE, "Using the Gaussian Image to Find the Orientation of Objects," *The International Journal of Robotics Research* 3(4) pp. 89-125 (Winter 1984).
5. CAYLEY, ARTHUR, "On certain results relating to quaternions," *Philosophical Magazine* xxvi pp. 141-145 (February 1845).
6. COURANT, R. AND HILBERT, D., *Methods of Mathematical Physics, Volume I*, Interscience Publishers, Inc., New York (1953).
7. DAHLQUIST, GERMUND AND BJÖRCK, ÅKE, *Numerical Methods*, Prentice-Hall, Inc., Englewood Cliffs, N.J. (1974). Translated by Ned Anderson.
8. EULER, LEONHARD, "Decouverte d'un nouveau principe de mécanique (1752)," pp. 81-108 in *Opera omnia, Ser. secunda, v. 5*, Orell Füssli Turici, Lausannae (1957).
9. EULER, LEONHARD, "Du mouvement de rotation des corps solides autour d'un axe variable (1758)," in *Opera omnia, Ser. secunda, v. 8*, Orell Füssli Turici, Lausannae ().
10. GABRIEL, STEVEN A. AND KAJIYA, JAMES T., "Spline Interpolation in Curved Manifolds," (1985). Submitted
11. GARDNER, MARTIN, *New Mathematical Diversions from Scientific American*, Fireside, St. Louis, Missouri (1971). Chapter 2
12. GOLDSTEIN, HERBERT, *Classical Mechanics, second edition*, Addison-Wesley Publishing Company, Inc., Reading, Mass. (1980). Chapter 4 and Appendix B.
13. GORDON, WILLIAM J. AND RIESENFELD, RICHARD F., "Bernstein-Bézier methods for the computer-aided design of free-form curves and surfaces," *J. ACM* 21(2) pp. 293-310 (April 1974).
14. GORDON, WILLIAM J. AND RIESENFELD, RICHARD F., "B-spline curves and surfaces," in *Computer Aided Geometric Design*, ed. Robert E. Barnhill and Richard F. Riesenfeld, Academic Press, New York (1974).
15. HAMILTON, SIR WILLIAM ROWAN, "On quaternions; or on a new system of imaginaries in algebra," *Philosophical Magazine* xxv pp. 10-13 (July 1844).
16. HERSTEIN, I.N., *Topics in Algebra, second edition*, John Wiley and Sons, Inc., New York (1975).
17. KANE, THOMAS R., LIKINS, PETER W. AND LEVINSON, DAVID A., *Spacecraft Dynamics*, McGraw-Hill, Inc. (1983).
18. LANE, JEFFREY M., CARPENTER, LOREN C., WHITED, TURNER, AND BLINN, JAMES F., "Scan line methods for displaying parametrically defined surfaces," *Comm. ACM* 23(1) pp. 23-34 (January 1980).
19. MACLANE, SAUNDERS AND BIRKHOFF, GARRETT, *Algebra, second edition*, Macmillan Publishing Co., Inc., New York (1979).
20. MISNER, CHARLES W., THORNE, KIP S., AND WHEELER, JOHN ARCHIBALD, *Gravitation*, W.H. Freeman and Company, San Francisco (1973). Chapter 41 — Spinors.
21. MITCHELL, E.E.L. AND ROGERS, A.E., "Quaternion Parameters in the Simulation of a Spinning Rigid Body," in *Simulation The Dynamic Modeling of Ideas and Systems with Computers*, ed. John McLeod, P.E., (1968).
22. NEWMAN, WILLIAM M. AND SPROULL, ROBERT F., *Principles of Interactive Computer Graphics, second edition*, McGraw-Hill, Inc., New York (1979). Chapter 21 — Curves and surfaces.
23. PICKERT, G. AND STEINER, H.-G., "Chapter 8 — Complex numbers and quaternions," in *Fundamentals of Mathematics, Volume I — Foundations of Mathematics: The Real Number System and Algebra*, ed. H. Behnke, F. Bachmann, K. Fladt, and W. Süss, (1983). Translated by S.H. Gould.
24. SCHMEIDLER, W. AND DREITZ, W., "Chapter 11 — Functional analysis," in *Fundamentals of Mathematics, Volume III — Analysis*, ed. H. Behnke, F. Bachmann, K. Fladt, and W. Süss, MIT Press, Cambridge, Mass. (1983). Translated by S.H. Gould.
25. SCHOENBERG, I.J., "Contributions to the problem of approximation of equidistant data by analytic functions," *Quart. Appl. Math.* 4 pp. 45-99 and 112-141 (1946).
26. SMITH, ALVY RAY, "Spline tutorial notes," Technical Memo No. 77, Computer Graphics Project, Lucasfilm Ltd. (May 1983).
27. SÜSS, W., GERICKE, H., AND BERGER, K.H., "Chapter 14 — Differential geometry of curves and surfaces," in *Fundamentals of Mathematics, Volume II — Geometry*, ed. H. Behnke, F. Bachmann, K. Fladt, and W. Süss, MIT Press (1983). Translated by S.H. Gould.
28. TAYLOR, RUSSELL H., "Planning and Execution of Straight Line Manipulator Trajectories," in *Robot Motion: Planning and Control*, ed. Michael Brady, John M. Hollerbach, Timothy L. Hohnson, Tomas Lozano-Perez and Matthew T. Mason, The MIT Press (1982).

Appendix I—Conversions

I.1 Quaternion to matrix

Using the restriction that $w^2 + x^2 + y^2 + z^2 = 1$ for a quaternion $q = [w, (x, y, z)]$, the formula for the corresponding matrix is

$$M = \begin{pmatrix} 1-2y^2-2z^2 & 2xy+2wz & 2xz-2wy \\ 2xy-2wz & 1-2x^2-2z^2 & 2yz+2wx \\ 2xz+2wy & 2yz-2wx & 1-2x^2-2y^2 \end{pmatrix}$$

If the quaternion does not have unit magnitude, an additional 4 multiplies and divides, 3 adds, and a square root will normalize it. (For the matrix conversion, the square root can be avoided in favor of divides if desirable.) Now we can obtain the operation count for creating the matrix. Most terms of the entries are a product of two factors, one of which is doubled. So we proceed as follows. First double x , y , and z , and form their products with w , x , y , and z . That will take 3 adds and 9 multiplies. Then form the sum for each of the 9 entries using 1 add each, plus an extra add for each of the 3 diagonal elements, for a total of 12 adds. Thus 9 multiplies and 15 adds suffice to convert a unit quaternion to a matrix.

I.2 Matrix to quaternion

An efficient way to determine quaternion components w , x , y , z from a matrix is to use linear combinations of the entries M_{mn} . Notice that the diagonal entries are formed from the squares of the quaternion components, while off-diagonal entries are the sum of a symmetric and a skew-symmetric part. Thus linear combinations of the diagonal entries will isolate squares of components; sums and differences of opposite off-diagonal entries will isolate products among x , y , and z and products with w . Using off-diagonals risks dividing by a component that may be zero, or within ϵ (the machine precision) of zero. However we can avoid that pitfall, and easily compute all components as follows.

$$w^2 = 1/4 (1 + M_{11} + M_{22} + M_{33})$$

$w^2 > \epsilon$?	
TRUE	FALSE
$w = \sqrt{w^2}$	$w = 0$
$x = (M_{23} - M_{32}) / 4w$	$x^2 = -1/2 (M_{22} + M_{33})$
$y = (M_{31} - M_{13}) / 4w$	$x^2 > \epsilon$?
$z = (M_{12} - M_{21}) / 4w$	
	TRUE
	FALSE
	$x = \sqrt{x^2}$
	$x = 0$
	$y = M_{12} / 2x$
	$y^2 = 1/2 (1 - M_{33})$
	$z = M_{13} / 2x$
	$y^2 > \epsilon$?
	TRUE
	FALSE
	$y = \sqrt{y^2}$
	$y = 0$
	$z = M_{23} / 2y$
	$z = 1$

No more than one square root, three divides, and a few adds and binary scales are required for any conversion.

I.3 Euler angles to quaternion

There are twelve possible axis conventions for Euler angles. The one used here is *roll*, *pitch*, and *yaw*, as used in aeronautics. A general rotation is obtained by first yawing around the z axis by an angle of ϕ , then pitching around the y axis by θ , and finally rolling around the x axis by ψ . Using the way quaternion components describe a rotation, we first obtain a quaternion for each simple rotation.

$$q_{roll} = [\cos \frac{\psi}{2}, (\sin \frac{\psi}{2}, 0, 0)]$$

$$q_{pitch} = [\cos \frac{\theta}{2}, (0, \sin \frac{\theta}{2}, 0)]$$

$$q_{yaw} = [\cos \frac{\phi}{2}, (0, 0, \sin \frac{\phi}{2})]$$

Multiplying these together in the right order gives the desired quaternion $q = q_{yaw} q_{pitch} q_{roll}$, with components

$$w = \cos \frac{\psi}{2} \cos \frac{\theta}{2} \cos \frac{\phi}{2} + \sin \frac{\psi}{2} \sin \frac{\theta}{2} \sin \frac{\phi}{2}$$

$$x = \sin \frac{\psi}{2} \cos \frac{\theta}{2} \cos \frac{\phi}{2} - \cos \frac{\psi}{2} \sin \frac{\theta}{2} \sin \frac{\phi}{2}$$

$$y = \cos \frac{\psi}{2} \sin \frac{\theta}{2} \cos \frac{\phi}{2} + \sin \frac{\psi}{2} \cos \frac{\theta}{2} \sin \frac{\phi}{2}$$

$$z = \cos \frac{\psi}{2} \cos \frac{\theta}{2} \sin \frac{\phi}{2} - \sin \frac{\psi}{2} \sin \frac{\theta}{2} \cos \frac{\phi}{2}$$

I.4 Euler angles to matrix

Combining the results of the previous two conversions gives

$$M =$$

$$\begin{pmatrix} \cos \theta \cos \phi & \cos \theta \sin \phi & -\sin \theta \\ \sin \psi \sin \theta \cos \phi - \cos \psi \sin \phi & \sin \psi \sin \theta \sin \phi + \cos \psi \cos \phi & \cos \psi \sin \theta \\ \cos \psi \sin \theta \cos \phi + \sin \psi \sin \phi & \cos \psi \sin \theta \sin \phi - \sin \psi \cos \phi & \cos \theta \cos \psi \end{pmatrix}$$

where ψ , θ , and ϕ are the angles of roll, pitch, and yaw, respectively.

I.5 Matrix to Euler angles

While converting a matrix to a unit quaternion only involves the sign ambiguity of square roots, converting to Euler angles involves inverse trigonometric functions, as we can only directly determine the sin's and cos's of the angles. Some convention, such as principle angles, must be adopted. However interpolation paths will vary greatly, depending on choice of angles. Setting that problem aside, here's a way to extract the sin's and cos's. Looking at the previous equation, $\sin \theta$ can be read off directly as $-M_{13}$. Use the trigonometric identity $\cos \theta = \pm \sqrt{1 - \sin^2 \theta}$ to compute $\cos \theta$ to within a sign, which is the best we can do. Assuming $\cos \theta$ is not zero, obtain the sin's and cos's of the other angles from

$$\sin \theta = -M_{13}$$

$$\cos \theta = \sqrt{1 - \sin^2 \theta}$$

$$\sin \psi = M_{23} / \cos \theta$$

$$\cos \psi = M_{33} / \cos \theta$$

$$\sin \phi = M_{12} / \cos \theta$$

$$\cos \phi = M_{11} / \cos \theta$$

If $\cos \theta$ is zero, then we must avoid dividing by zero. It also becomes impossible to distinguish roll from yaw. Adopting the convention that the yaw angle ϕ is 0 allows

$$\sin \psi = -M_{32}$$

$$\cos \psi = M_{22}$$

$$\sin \phi = 0$$

$$\cos \phi = 1$$

From these values a two argument \tan^{-1} will give angles between $-\pi$ and $+\pi$, or 0 and 2π , or some other conventional range; take your pick. (For a faster conversion, just compute, say, \sin^{-1} and check the sign of the cosine term with respect to $\cos \theta$.) Because of the uncertainties of square roots, inverse trigonometric functions, and yaw-roll separation, matrix to Euler angle conversion is inherently *very* ill-defined.

1.6 Quaternion to Euler angles

Use the most straight-forward approach: convert the quaternion to a matrix, then the matrix to Euler angles. Of course it is unnecessary to compute matrix elements that are never used. This conversion is also unavoidably ill-defined, as quaternions contain no more information about angles than matrices do.